

LETTER TO THE EDITOR

Severe chronic psychosis after allogeneic SCT from a schizophrenic sibling

Bone Marrow Transplantation (2015) **50**, 153–154; doi:10.1038/bmt.2014.221; published online 6 October 2014

Schizophrenia is a life-long disorder, usually starting at early adulthood and consisting of remitting or chronic psychosis and functional decline.

There is ample evidence that immune processes play a role in schizophrenia.^{1–4} More than 20 different auto-antibodies are at elevated levels in patients with schizophrenia as compared to controls.⁵ Autoimmune diseases (AID), such as thyrotoxicosis, celiac disease, acquired hemolytic anaemia, interstitial cystitis and Sjögren's syndrome, but also atopic diseases, have higher prevalence rates among patients with schizophrenia.^{6,7} The strongest genetic association with schizophrenia is found in the MHC genes, including loci that influence immune responses.⁸ Further evidence for an autoimmune component comes from the lower incidence of schizophrenia in men who used glucocorticosteroids for somatic diseases (odds ratio 0.52).⁹ It is unclear if immune processes play a key role in all patients with schizophrenia, or only in ~30–40% of them.^{10,11}

A common characteristic of AID¹² and severe allergies¹³ is their favourable response to immune ablation and rescue with SCT. Accordingly, we (IES and DWvB) searched for transplant cases suffering from coincident schizophrenia.^{14,15} No such cases had been discovered, but one of us (GT) retrieved the history of a patient who developed severe psychosis after receiving a BM transplant from his schizophrenic brother.

The patient (born 1940) had a blank psychiatric history. He was retired, married and had two adult children. At the age of 67 he developed fatigue and skin ecchymoses, and he was diagnosed with chronic lymphocytic leukemia (CLL) and bone marrow aplasia, requiring weekly blood transfusions. Chemotherapy (two courses of CY), and treatment with cyclosporin A, rituximab and prednisolone did not improve his CLL/aplastic anemia. In 2007 he received an allogeneic peripheral blood SCT from one of his brothers, who was the only HLA-matched family member. This brother (born 1952) had schizophrenia since early adulthood and he required treatment with multiple antipsychotic agents. No other first-degree relatives suffered from schizophrenia. The donor also had a history of Lyme disease 8 years before. Serology showed positive IgG and negative IgM for Lyme disease, and PCR was negative for *Borrelia burgdorferi* DNA, consistent with inactive prior infection. The donor used doxycycline 100 mg twice daily, starting 4 days before the stem cell collection, as a safety measure to prevent transmission. The patient was conditioned with fludarabine and CY, followed by an infusion of 5.0×10^6 CD34⁺ cells/kg from the peripheral blood of his brother. Tacrolimus was administered to prevent GVHD. He never received steroids, and did not develop graft-vs-host reactions. He had complete hematologic recovery and reached full hematopoietic chimerism (>97% donor cells 4 weeks after SCT). We tapered off tacrolimus 4 months after the SCT, because of decreasing blood counts. A few weeks later, when off tacrolimus, he developed acute psychotic symptoms: frequent hallucinations (running commentary and threatening voices), bizarre and non-bizarre delusions, and thought broadcasting with clear consciousness. Insight and

judgment were poor. He described his mood as 'angry', with a flat affect. When he developed suicidal and homicidal ideation, he was admitted to a psychiatric clinic. Neurological evaluation, magnetic resonance imaging and electroencephalography revealed no abnormalities. Extensive medical work-up did not reveal metabolic disorders, underlying infections and neoplasms. Results of lumbar puncture were normal, and screenings for viruses (including Herpes Simplex Virus and Human Herpes Virus-6), bacteria, fungi and Lyme disease were all unremarkable. He was treated with risperidone 3 mg and citalopram 20 mg, unsuccessfully. Under the working diagnosis of delirium, all medications were discontinued and haloperidol 1 mg was administered, which was not helpful either.

The patient's family decided for comfort care only and the patient was lost to follow-up. He died in 2010, of unknown cause.

The following differential diagnoses were considered and dismissed:

Delirium: Acute onset of psychotic symptoms in a 68-year-old man after somatic disease. However, the 4-month interval between transplantation and onset of psychosis is atypical. Furthermore, discontinuation of all medication did not improve his condition, nor could any somatic disorder be identified.

Endogenous schizophrenia: The patient was genetically predisposed to schizophrenia, given his brother had this disease. However, acute onset at the age of 68 is rare,¹⁶ especially in males.¹⁷ Age of onset is strongly correlated among affected siblings, and differences in onset of more than 10 years are very rare.¹⁸

Tick-borne infection transmitted through stem cells: The stem cell donor had Lyme disease 8 years earlier. Transfer of tick-borne pathogens might have caused psychotic symptoms. However, the donor was *Borrelia* IgM and PCR negative and received doxycycline before stem cell extraction.

Therefore, adoptive transfer of schizophrenia is the most appealing etiology, not only by exclusion, but also in view of the increasing evidence that some forms of schizophrenia have an autoimmune origin. Although the transplanted patient fulfills these criteria, a formal diagnosis was never made given his very unusual age of onset and potential relation to the SCT.

Adoptive transfer of AIDs seems to be rare, as we are aware of only 21 cases reported so far among ~200 000 long-term survivors of allogeneic SCTs. Stem cell donors are routinely subjected to complete medical examination; therefore chances are small that AID is overlooked. The 21 cases include thyroiditis (10 cases),¹⁹ vitiligo (3),^{20–22} psoriasis (2),^{23,24} type I diabetes mellitus (2),²⁵ celiac disease (1),²⁶ thrombocytopenia (1),²⁷ polyglandular syndrome type II (1)²² and systemic lupus erythematosus (1).²⁸ Moreover, not only AIDs may be transferred by SCT but also other immune diseases, such as allergies.¹³ Adoptive transfer of AIDs and allergies is thought to be mediated by transfer of donor lymphocytes. This suggests that the subform of schizophrenia in the patient was mediated by lymphocytes.

Based on this single case report, we obviously cannot prove an immune pathogenesis of schizophrenia. However, the report supports the hypothesis of immunological involvement in schizophrenia pathogenesis and we suggest that physicians and patients involved in SCT take into consideration the possibility that schizophrenia may be transmitted by the transplant.

Taking it one step further, coexisting schizophrenia might not *per se* be considered as a contraindication in patients with standard transplant indications, but rather as an opportunity for inducing remission of the psychotic process and as a potential area of research.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

This work is supported by a grant from the Dutch Scientific Council (ZonMw) (TOP grant Microglia cells in schizophrenia: a potential new target for treatment).

IE Sommer¹, DW van Bekkum², H Klein³, R Yolken⁴, Lot de Witte¹ and G Talamo⁵

¹Department of Psychiatry, University Medical Center Utrecht, Utrecht, The Netherlands;

²Department of Hematology, Erasmus University Medical Center, Rotterdam, The Netherlands;

³Department of Psychiatry, Groningen University, University Medical Centre Groningen, Groningen, The Netherlands;

⁴Department of Pediatrics, Stanley Division of Developmental Neurovirology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD, USA and

⁵Bone Marrow Transplant Penn State Hershey Cancer Institute Hershey, Hershey, PA, USA
E-mail: dw@bekkum.nu

REFERENCES

- Benros ME, Mortensen PB, Eaton WW. Autoimmune diseases and infections as risk factors for schizophrenia. *Ann N Y Acad Sci* 2012; **1262**: 56–66.
- Goldsmith CA, Rogers DP. The case for autoimmunity in the etiology of schizophrenia. *Pharmacotherapy* 2008; **28**: 730–741.
- Jones AL. Immune dysregulation and self-reactivity in schizophrenia: Do some cases of schizophrenia have an autoimmune basis? *Immunol Cell Biol* 2005; **83**: 9–17.
- Gibney SM, Drexhage HA. Evidence for a dysregulated immune system in the etiology of psychiatric disorders. *J Neuroimmune Pharmacol* 2013; **8**: 900–920.
- Ezeoke A, Mellor A, Buckley P, Miller B. A systematic, quantitative review of blood autoantibodies in schizophrenia. *Schizophr Res* 2013; **150**: 245–251.
- Eaton WW, Byrne M, Ewald H, Mors O, Chen CY, Agerbo E *et al.* Association of schizophrenia and autoimmune diseases: linkage of Danish national registers. *Am J Psychiatry* 2006; **163**: 521–528.
- Pedersen MS, Benros ME, Agerbo E, Børglum AD, Mortensen PB. Schizophrenia in patients with atopic disorders with particular emphasis on asthma: a Danish population-based study. *Schizophr Res* 2012; **138**: 58–62.
- De Jong S, van Eijk KR, Zeegers DW, Strengman E, Janson E, Veldink JH *et al.* Expression QTL analysis of top loci from GWAS meta-analysis highlights additional schizophrenia candidate genes. *Eur J Hum Genet* 2012; **20**: 1004–1008.
- Laan W, Smeets H, de Wit NJ, Kahn RS, Grobbee DE, Burger H. Glucocorticosteroids associated with a decreased risk of psychosis. *J Clin Psychopharmacol* 2009; **29**: 288–290.
- Schwarz E, van Beveren NJ, Ramsey J, Leweke FM, Rothermundt M, Bogerts B *et al.* Identification of subgroups of schizophrenia patients with changes in either immune or growth factor and hormonal pathways. *Schizophr Bull* 2013; **40**: 787–795.
- Fillman SG, Cloonan N, Miller LC, Weickert CS. Markers of inflammation in the prefrontal cortex of individuals with schizophrenia. *Mol Psychiatry* 2013; **18**: 133.
- Tyndall A. Successes and failures of stem cell transplantation in autoimmune diseases. *Hematology Am Soc Hematol Educ Program* 2011; **2011**: 280–284.
- Khan F, Hallstrand TS, Geddes MN, Henderson WR Jr, Storek J. Is allergic disease curable or transferable with allogeneic hematopoietic cell transplantation? *Blood* 2009; **113**: 279–290.
- Sommer IE, van Bekkum DW. Call for case histories of BMT in patients with coincident schizophrenia. *Leukemia* 2013; **27**: 1217–1218.
- Sommer IE, van Bekkum DW. Call for case histories of BMT in patients with coincident schizophrenia. *Bone Marrow Transplant* 2013; **48**: 880.
- Girard C, Simard M. Clinical characterization of late- and very late-onset first psychotic episode in psychiatric inpatients. *Am J Geriatr Psychiatry* 2008; **16**: 478–487.
- Meesters PD, de Haan L, Comijs HC, Stek ML, Smeets-Janssen MM, Weeda MR *et al.* Schizophrenia spectrum disorders in later life: prevalence and distribution of age at onset and sex in a Dutch catchment area. *Am J Geriatr Psychiatry* 2012; **20**: 18–28.
- Burke JG, Murphy BM, Bray JC, Walsh D, Kendler KS. Clinical similarities in siblings with schizophrenia. *Am J Med Genet* 1996; **67**: 239–243.
- Marazuela M, Steegman JL. Transfer of autoimmune hypothyroidism following bone marrow transplantation from a donor with Graves' disease. *Bone Marrow Transplant* 2000; **26**: 1217–1220.
- Alajlan A, Alfadley A, Pedersen KT. Transfer of vitiligo after allogeneic bone marrow transplantation. *J Am Acad Dermatol* 2002; **46**: 606–610.
- Neumeister P, Strunk D, Apfelbeck U, Sill H, Linkesch W. Adoptive transfer of vitiligo after allogeneic bone marrow transplantation for non-Hodgkin's Lymphoma. *Lancet* 2000; **355**: 1334–1335.
- Mellouli F, Ksouri H, Lakhal A, Torjmen L, Ladeb S, Ben Othman T *et al.* Autoimmune polyglandular syndrome type II after bone marrow transplant: real transfer or acceleration of a programmed disease? *Exp Clin Transplant* 2012; **10**: 76–80.
- Gardembas-Pain M, Ifrah N, Foussard C, Boasson M, Andre JPS, Verret JL. Psoriasis after allogeneic bone marrow transplantation [letter]. *Arch Dermatol* 1990; **126**: 1523.
- Snowden JA, Heaton DC. Development of psoriasis after syngeneic bone marrow transplant from psoriatic donor: further evidence for adoptive autoimmunity. *Br J Dermatol* 1997; **137**: 130–132.
- Lampeter EF, McCann SR, Kolb H. Transfer of diabetes type 1 by bone-marrow transplantation. *Lancet* 1998; **351**: 568–569.
- Bargetzi MI. Celiac disease transmitted by allogeneic non-T cell depleted bone marrow transplantation. *Bone Marrow Transplant* 1997; **20**: 607–609.
- Waters AH, Metcalfe P, Minchinton RM, Barrett AJ, James DC. Autoimmune thrombocytopenia acquired from allogeneic bone marrow graft: compensated thrombocytopenia in bone marrow donor and recipient. *Lancet* 1983; **2**: 1430.
- Ritchie DS, Sainani A, D'Souza A, Grigg AP. Passive donor-to-recipient transfer of antiphospholipid syndrome following allogeneic stem-cell transplantation. *Am J Hematol* 2005; **79**: 299–302.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>